VOL. 692, NOS. 1 + 2 JOURNAL OF CHROMATOGRAPHY A 10 FEBRUARY 1995

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index)

18TH INTERNATIONAL SYMPOSIUM ON COLUMN LIQUID CHROMATOGRAPHY, MINNEAPOLIS, MN, USA, 8–13 MAY 1994, PART II

ENVIRONMENTAL APPLICATIONS

Automated procedure for determination of trace amounts of aldehydes in drinking water
by R. Wu and L.B. White (Fremont, CA, USA) 1

Automated precolumn concentration and high-performance liquid chromatographic analysis of polynuclear aromatic hydrocarbons in water using a single pump and a single valve
by F. Lai and L. White (Fremont, CA, USA) 11

Optimization of automated solid-phase extraction for quantitation of polycyclic aromatic hydrocarbons in aqueous media by high-performance liquid chromatography–UV detection
by N.C. Fladung (Johnston, IA, USA) 21

High-performance liquid chromatographic determination of sulfonylureas in soil and water
by G.C. Galletti (Gallina, Italy) and A. Bonetti and G. Dinelli (Bologna, Italy) 27

Liquid chromatographic determination of the mycotoxin fumonisin B_2 in physiological samples
by G.S. Shephard, P.G. Thiel and E.W. Sydenham (Tygerberg, South Africa) 39

PHARMACEUTICALS, DRUGS, BIOMEDICAL APPLICATIONS

Determination of L-735 524, an human immunodeficiency virus protease inhibitor, in human plasma and urine via high-performance liquid chromatography with column switching
by E. Woolf, T. Au, H. Haddix and B. Matuszewski (West Point, PA, USA) 45

Comparison of high-performance liquid chromatographic methods for the determination of 1,2-dibromo-2,4-dicyanobutane in cosmetic products
by S.C. Rastogi and S.S. Johansen (Roskilde, Denmark) 53

Alkylsulphonic acid ion pairing with radial compression columns for determining plasma or cerebrospinal fluid 1-β-D-arabinofuranosylcytosine in pediatric pharmacokinetic analysis
by M.L. Stout and Y. Ravindranath (Detroit, MI, USA) 59

High-performance liquid chromatographic method for potency determination of cephalexin in commercial preparations and for stability studies
by M.-C. Hsu (Taoyuan, Taiwan) and Y.-S. Lin and H.-C. Chung (Taipei, Taiwan) 67

Rapid liquid chromatographic–mass spectrometric assay for oxymetazoline in whole rat blood
by F.J. Hayes, T.R. Baker, R.L.M. Dobson and M.S. Tsueda (Cincinnati, OH, USA) 73

Hydrophobicity parameter from high-performance liquid chromatography on an immobilized artificial membrane column and its relationship to bioactivity
by A. Nasal, M. Sznitowska, A. Buciński and R. Kaliszan (Gdańsk, Poland) 83

Unusual examples of the liquid chromatographic resolution of racemates. Resolution of π-donor analytes on a π-donor chiral stationary phase
by M.H. Hyun, J.-J. Ryoo, Y.J. Cho and J.S. Jin (Pusan, South Korea) 91

High-performance liquid chromatographic determination of (4-[^{11}C]methoxyphenyl)-(5-fluoro-2-hydroxyphenyl)-methyleneaminobutyric acid and its benzophenone metabolite
by F. De Vos and G. Slegers (Ghent, Belgium) 97

Use of high-performance liquid chromatography–diode array detection in forensic toxicology
by E.M. Koves (Toronto, Canada) 103

Qualitative and quantitative determination of illicit heroin street samples by reversed-phase high-performance liquid chromatography: method development by CARTAGO-S
by K. Grogg-Sulser, H.-J. Helmlin and J.-T. Clerc (Berne, Switzerland) 121

Determination of aflatoxins in medicinal herbs and plant extracts
by K. Reif and W. Metzger (Verstenbergsgreuth, Germany) 131

Determination of liquiritin, glycyrrhizin, hesperidin, cinnamic acid, cinnamaldehyde, honokiol and magnolol in the traditional Chinese medicinal preparation Wuu-Ji-San by high-performance liquid chromatography
by Y.-C. Lee, C.-Y. Huang, K.-C. Wen and T.-T. Suen (Taipei, Taiwan) 137

Chemometric categorization of octadecylsilyl bonded-phase silica columns using test mixtures and confirmation of results with pharmaceutical compound separations
by B.A. Olsen and G.R. Sullivan (Lafayette, IN, USA) 147

Liquid chromatographic determination of the macrolide antibiotics roxithromycin and clarithromycin in plasma by automated solid-phase extraction and electrochemical detection
by M. Hedenmo and B.-M. Eriksson (Mölndal, Sweden) 161

Analysis of mycolic acids by high-performance liquid chromatography and fluorimetric detection. Implications for the identification of mycobacteria in clinical samples
by S.R. Hagen and J.D. Thompson (Hamilton, MT, USA) 167

Resolution of promethazine, ethopropazine, trimeprazine and trimipramine enantiomers on selected chiral stationary phases using high-performance liquid chromatography
by G.W. Ponder (Athens, GA, USA), S.L. Butram (Dahlonega, GA, USA), A.G. Adams (Marietta, GA, USA) and C.S. Ramanathan and J.T. Stewart (Athens, GA, USA) 173

Comparison of theory-based and empirical modeling for the prediction of chromatographic behavior in the ion-pairing separation of benzodiazepine-derived pharmaceutical compounds
by L.A. Larew, B.A. Olsen, J.D. Stafford and M.V. Wilhelm (Lafayette, IN, USA) 183

Simultaneous determination of granisetron and its 7-hydroxy metabolite in human plasma by reversed-phase high-performance liquid chromatography utilizing fluorescence and electrochemical detection
by V.K. Boppana (King of Prussia, PA, USA) 195

Determination of citrate, inositol and gentisic acid in a pharmaceutical diagnostic formulation by ion-moderated partition chromatography (Short communication)
by R.W. Chong and B.J. Moore (St. Louis, MO, USA) 203

Use of overlapping resolution mapping scheme for optimization of the high-performance liquid chromatographic separation of pharmaceuticals
by C.P. Ong, K.K. Chow, C.L. Ng, F.M. Ong, H.K. Lee and S.F.Y. Li (Singapore, Singapore) 207

APPLICATIONS, OTHERS

Preparative high-performance liquid chromatography for the purification of natural anthocyanins
by M. Fiorini (Bologna, Italy) 213

Compositional distribution characterization of poly(methyl methacrylate)–graft-polydimethylsiloxane copolymers
by T.C. Schunk (Rochester, NY, USA) and T.E. Long (Kingsport, TN, USA) 221

Rapid scaleable chromatographic purification of nucleic acids from proteinaceous mixtures (Short communication)
by Y.L. Hofman (Wilmington, DE, USA), K.L. Petrillo (Newark, DE, USA), H.C. Greenblatt and R. Lehrer (Wilmington, DE, USA) and M.S. Payne (Newark, DE, USA) 233

Separation of photosynthetic pigments and their precursors by reversed-phase high-performance liquid chromatography using a photodiode-array detector
by B. Schoefs (Villeneuve d'Ascq, France), M. Bertrand (Tourlaville, France) and Y. Lemoine (Villeneuve d'Ascq, France) 239

Analysis of nerve agent degradation products using capillary ion electrophoresis (Short communication)
by S.A. Oehrle (Milford, MA, USA) and P.C. Bossle (Aberdeen Proving Ground, MD, USA) 247

Extending the scope of chiral separation of basic compounds by cyclodextrin-mediated capillary zone electrophoresis
by C. Quang and M.G. Khaledi (Raleigh, NC, USA) 253

Determination of polycarboxylic acids by capillary electrophoresis with copper complexation
by J.P. Wiley (Edgewater, NJ, USA) 267

Mass spectrometric detectors for samples separated by planar electrophoresis (Review)
by K.L. Busch (Atlanta, GA, USA) 275

Selectivity control in micellar electrokinetic chromatography of small peptides using mixed fluorocarbon–hydrocarbon anionic surfactants
by B. Ye, M. Hadjmohammadi and M.G. Khaledi (Raleigh, NC, USA) 291

Linear solvation energy relationships in micellar liquid chromatography and micellar electrokinetic capillary chromatography
by S. Yang and M.G. Khaledi (Raleigh, NC, USA) 301

Micellar liquid chromatographic separation of sulfonamides in physiological samples using direct on-column injection
by S. Yang and M.G. Khaledi (Raleigh, NC, USA) 311

Experimental evidence for the existence of duoselective (Type III) enantiomer separations in the capillary electrophoretic analysis of chiral weak acids
by M.E. Biggin, R.L. Williams and G. Vigh (College Station, TX, USA) 319

AUTHOR INDEX VOLS. 691 AND 692 327